



# Evolution of the HER/ErbB Receptor System

Worms:	1 Ligand and 1 Receptor
Flies:	4 Ligands and 1 Receptor
Mammals:	12 Ligands and 4 Receptors

# The HERs A Dysfunctional Family of Receptors






# HER2 Activation in Cancer

Ligand-independent  
(Amplified HER2 tumors)

Ligand-dependent  
(Non-amplified HER2 tumors)



Herceptin patient population

# HER2 Associates with HER3 in a Heregulin-Dependent Manner

- 2C4 blocks ligand-dependent HER2-HER3 association, Herceptin does not.

IP:  $\alpha$ HER2

MCF7


Low/Normal ErbB2



WB:  $\alpha$ HER3

SK-BR-3

High ErbB2



83 -

83 -





HRG:	-	+	-	+	-	+	-	+	-	+	-	+	-	+	-	+
	1	2	3	4	1	2	3	4	1	2	3	4	1	2	3	4

1. Control    2. 2C4    3. Herceptin    4.  $\alpha$ EGFR





Rob Akita

# Ovarian Tumor Cell Lines

	<u>3</u>		<u>420</u>		<u>429</u>		<u>432</u>	
2C4	-	+	-	+	-	+	-	+
HRG	-	+	-	+	-	+	-	+




IP: H2  
WB: H3



IP: H2  
WB: H2

# ErbB2 is recruited to ErbB3-HRG Complexes



ErbBX

# 2C4 Disrupts Ligand-Dependent ErbB2 Signaling

32



2C4Y

2C4Y

Low  
Affinity  
Receptor

High  
Affinity  
Receptor

Ligand-Activated  
Hetero-oligomer



ATP  
ADP


ErbBX




## EGFR

## EGFR-EGF Complex

Closed



Open




*Ogiso et al. Cell (2002) 110: 775*


*Garret et al. Cell (2002) 110: 763*

*Ferguson et al. Mol Cell (2003) 11:507*

# EGFR-EGF Complex




# HER2



*Cho et al. Nature 421:756.  
Matt Franklin & Bart de Vos, Genentech*

# Receptor dimerization via a domain II handshake



# Comparison of HER2-EGFR to EGFR-EGFR Complexes

Characteristics of heterocomplexes :

Higher affinity

Decreased internalization rates

Altered trafficking

Diverse downstream signaling

# HER2-HER3 Complex: A Paradigm for Efficient Molecular Collaboration

Symbiotic relationship

Ligand-less HER2 and defective-kinase HER3  
Most potent HER signaling complex.  
Efficiently activates both MAPK and PI3K signaling pathways.


HER2's active kinase

HER3 serves as a kinase substrate for HER2.

Multiple potential tyrosine phosphorylation sites.  
Especially for PI3-kinase.


Most active complex with regard to transformation potential.

# Heregulin Binding to Cos Cell Transfectants




Gabriele Schaefer

# Scatchard Analysis of Heregulin Binding to Cells Transfected with HER3 and HER2




*Gabriele Schaefer*

# Inhibition of Heregulin Binding by 2C4 a Monoclonal Antibody to ErbB2




Gabriele Schaefer







*Cho et al. Nature 421:756.*  
*Matt Franklin & Bart de Vos, Genentech*



# 2C4 Inhibits Heregulin-Dependent HER3-HER2 Signaling




# Coupling of HER2/3 to the MAPK and Akt Pathways




# 2C4 Inhibits Ligand-Dependent HER2 Signaling (MAPK)

**B**



# Heregulin-Dependent Akt Activation 2C4 Inhibits

ErbB3



GSK3  $\alpha/\beta$  →

HRG

2C4

Heregulin

-	+	-	+	-	+
-	-	+	+	-	-
-	-	-	-	+	+

1035

1178

1180

grb7

1203/05

1241

1243

grb7

1257


1270

1309

shc


Rob Akita

**HER2/HER3 receptor activation increases the rate of hypoxia-inducible factor (HIF-1 $\alpha$ ) synthesis**



# 2C4 Blocks Heregulin-Induced Expression of VEGF

Relative  
VEGF  
mRNA  
Expression





## Genentech Acknowledgments

Rob Akita

Gabriele Schaefer

Julie Lofgren

Paul Pisacane

Ralph Schwall

Lisa Crocker

Gail Phillips

Klara Totpol

Inessa Balter

Cam Adams

Len Presta

Matt Franklin

Ken Carey

Bart de Vos

# Prostate Cancer and HER2

- Clinical studies:

HER2 gene amplification or protein overexpression is rare.

HER ligand expression (e.g., TGF- $\alpha$ ) frequently occurs with the onset of the androgen-independent phenotype.

# Prostate Cancer and HER2

## • Laboratory studies:

Onset of the androgen-independent phenotype corresponds with HER2 overexpression. (Sawyers).


Evidence for cross-talk between HER2 and androgen receptor signal transduction pathways (Chung).

# Androgen-Independent Prostate Cancer

Autocrine  
activation of HER-  
kinase axis

Dysregulation of  
AR; *unresponsive*  
to androgen  
ablation

Increased  
expression of  
HER2?



*adapted from Kim et al. (1999)*

# CWR Prostate Cancer Models

Derived from a primary prostate cancer patient by Thomas Pretlow, Case Western Reserve.

Xenograft maintained by serial transplantation in nude mice.


Growth is androgen-dependent (CWR22).

Good correlation between tumor growth and serum PSA levels.


Tumors regress after androgen withdraw.

Relapsed tumors are androgen-independent (CWR22R & CWRSA6).

# Relative Expression of ErbB Receptors and Ligands in CWR Tumors




# Effect of 2C4 on the Growth of the Androgen-Independent Human Prostate Cancer Xenograft CWRSA6



# Proof of Concept Experiment: 2C4 Does Not Require An Intact Fc For Anti-Tumor Activity

## Androgen-Independent Prostate Cancer Xenograft (CWR22R)






Breece & Schmelzer

Agus & Higgins, MSK



# Effect of 2C4 on the Growth of the Androgen-Dependent Human Prostate Cancer Models



# Summary of prostate cancer studies

In contrast to Herceptin<sup>®</sup>, 2C4 inhibits the growth of androgen-independent prostate tumor xenografts


model represents a patient population that is readily available for clinical studies

Combining 2C4 with low-dose Taxol<sup>®</sup> results in significant tumor regression and in many cases tumor elimination


2C4 also inhibits the growth of androgen-dependent prostate tumor xenografts. These data suggest that 2C4 may be active in patients with early-stage prostate cancer

# Breast Cancer Studies

# 2C4 Has Herceptin-Like Activity Against High HER2 Expressing Tumors




# Effect of rhuMAb 2C4 or Herceptin® on the growth of human breast cancer cells (low HER2 expression)



-3 day assay  
(Alamar Blue)

# Evaluation of rhuMAb 2C4 in the breast cancer xenograft MAXF 449 (low HER2 expression)


Treatment schedule: i.p.; once/week (Day 1, 8, 15, 22, 29 and 36; 2x loading dose at day 1)



HH Fiebig, Oncotest

# Properties of rhuMAb 2C4

## Trastuzumab Herceptin



- Binds in IV near JM.
- Protects against receptor shedding.
- Moderately affects receptor down-modulation.
- Slight effect on HER2's role as a coreceptor.

## Pertuzumab 2C4



- Binds in II at dimerization interface.
- Does not prevent receptor shedding.
- Moderately affects receptor down-modulation.
- Major effect on HER2's role as a coreceptor.



# Collaborators

David Agus: Cedars Sinai

Howard Scher: Memorial Sloan-Kettering

Hans-Joachim Mueller: Roche-Penzberg

HH Fiebig: Oncotest Freiburg